

APPROXIMATE SOLUTIONS TO VARIATIONAL INEQUALITY OVER THE FIXED POINT SET OF A STRONGLY NONEXPANSIVE MAPPING

SHIGERU IEMOTO, KAZUHIRO HISHINUMA, AND HIDEAKI IIDUKA

ABSTRACT. Variational inequality problems over fixed point sets of nonexpansive mappings include many practical problems in engineering and applied mathematics, and a number of iterative methods have been presented to solve them. In this paper, we discuss a variational inequality problem for a monotone, hemicontinuous operator over the fixed point set of a *strongly* nonexpansive mapping on a real Hilbert space. We then present an iterative algorithm, which uses the strongly nonexpansive mapping at each iteration, for solving the problem. We show that the algorithm potentially converges in the fixed point set faster than algorithms using firmly nonexpansive mappings. We also prove that, under certain assumptions, the algorithm with slowly diminishing step-size sequences converges to a solution to the problem in the sense of the weak topology of a Hilbert space. Numerical results demonstrate that the algorithm converges to a solution to a concrete variational inequality problem faster than the previous algorithm.

1. INTRODUCTION

The paper presents an iterative algorithm for the variational inequality problem [30, 32, 36, 38–40, 43, 44] for a monotone, hemicontinuous operator A over a nonempty, closed convex subset C of a real Hilbert space H with inner product $\langle \cdot, \cdot \rangle$ and its induced norm $\|\cdot\|$,

$$(1) \quad \text{find } z \in C \text{ such that } \langle y - z, Az \rangle \geq 0 \text{ for all } y \in C.$$

Problem (1) can be solved by using convex optimization techniques. A typical iterative procedure for Problem (1) is the *projected gradient method* [17, 43], and it is expressed as $x_1 \in C$ and $x_{n+1} = P_C(I - r_n A)x_n$ for $n = 1, 2, \dots$, where P_C stands for the metric projection onto C , I is the identity mapping on H , and $\{r_n\} \subset (0, \infty)$. However, as the method requires repetitive use of P_C , it can only be applied when the explicit form of P_C is known (e.g., C is a closed ball or a closed cone). The following method, called the *hybrid steepest descent method* (HSDM) [42], enables us to consider the case in which C has a more complicated form: $x_1 \in H$ and

$$x_{n+1} = (I - r_n A)Tx_n$$

for all $n = 1, 2, \dots$, where $\{r_n\} \subset (0, 1]$ and $T: H \rightarrow H$ is an easily implemented nonexpansive mapping satisfying $\text{Fix}(T) := \{x \in H: Tx = x\} = C$. HSDM

2000 *Mathematics Subject Classification.* 47H06, 47J20, 47J25.

Key words and phrases. variational inequality problem, fixed point set, strongly nonexpansive mapping, monotone operator

This work was supported by the Japan Society for the Promotion of Science through a Grant-in-Aid for JSPS Fellows (08J08592) and a Grant-in-Aid for Young Scientists (B) (23760077).

converges strongly to a unique solution to the variational inequality problem over $\text{Fix}(T)$,

$$(2) \quad \text{find } z \in \text{Fix}(T) \text{ such that } \langle y - z, Az \rangle \geq 0 \text{ for all } y \in \text{Fix}(T),$$

when $A: H \rightarrow H$ is strongly monotone and Lipschitz continuous. Problem (2) contains many applications such as signal recovery problems [13], beamforming problems [37], power control problems [22, 28], bandwidth allocation problems [23–25], and optimal control problems [29]. References [13], [26], and [27] presented acceleration methods for solving Problem (2) when A is strongly monotone and Lipschitz continuous. Algorithms were presented to solve Problem (2) when A is (strictly) monotone and Lipschitz continuous [23, 25]. When $H = \mathbb{R}^N$ and $A: \mathbb{R}^N \rightarrow \mathbb{R}^N$ is continuous (and is not necessarily monotone), a simple algorithm, $x_{n+1} := \alpha_n x_n + (1 - \alpha_n)(1/2)(I + T)(x_n - r_n A x_n)$ ($\alpha_n, r_n \in [0, 1]$), was presented in [22] and the algorithm converges to a solution to Problem (2) under some conditions.

Reference [20] proposed an iterative algorithm for solving Problem (2) when $A: H \rightarrow H$ is monotone and hemicontinuous and showed that the algorithm weakly converges to a solution to the problem under certain assumptions. The results in [20] are summarized as follows: suppose that $F: H \rightarrow H$ is a firmly nonexpansive mapping with $\text{Fix}(F) \neq \emptyset$ and that $A: H \rightarrow H$ is a monotone, hemicontinuous mapping with

$$\text{VI}(\text{Fix}(F), A) := \{z \in \text{Fix}(F) : \langle y - z, Az \rangle \geq 0 \text{ for all } y \in \text{Fix}(F)\} \neq \emptyset.$$

Define a sequence $\{x_n\} \subset H$ by $x_1 \in H$ and

$$(3) \quad \begin{cases} y_n = F(I - r_n A)x_n, \\ x_{n+1} = \alpha_n x_n + (1 - \alpha_n)y_n \end{cases}$$

for all $n = 1, 2, \dots$, where $\{\alpha_n\} \subset [0, 1)$ and $\{r_n\} \subset (0, 1)$. Assume that $\{Ax_n\}$ in Algorithm (3) is bounded, and that there exists $n_0 \in \mathbb{N}$ such that $\text{VI}(\text{Fix}(F), A) \subset \Omega := \bigcap_{n=n_0}^{\infty} \{x \in \text{Fix}(F) : \langle x_n - x, Ax_n \rangle \geq 0\}$. If $\{\alpha_n\}$ and $\{r_n\}$ satisfy $\limsup_{n \rightarrow \infty} \alpha_n < 1$, $\sum_{n=1}^{\infty} r_n^2 < \infty$, and $\lim_{n \rightarrow \infty} \|x_n - y_n\|/r_n = 0$, then $\{x_n\}$ weakly converges to a point in $\text{VI}(\text{Fix}(F), A)$. To relax the strong monotonicity condition of A considered in [42], a *firmly nonexpansive mapping* F is used in Algorithm (3) in place of a nonexpansive mapping T . From the fact that a firmly nonexpansive mapping F can be represented by the form, $F = (1/2)(I + T)$, for some nonexpansive mapping T , Algorithm (3) when $\alpha_n := 0$ and $F := (1/2)(I + T)$ can be simplified as follows: $x_1 \in H$ and

$$(4) \quad x_{n+1} = \frac{1}{2}(I + T)(x_n - r_n A x_n) = \frac{1}{2}(x_n - r_n A x_n) + \frac{1}{2}T(x_n - r_n A x_n).$$

In constrained optimization problems, one is required to satisfy constraint conditions early in the process of executing an iterative algorithm. From this viewpoint, we introduce the following algorithm with a weighted parameter α , which is more than $1/2$:

$$(5) \quad \begin{aligned} x_{n+1} &= (1 - \alpha)(x_n - r_n A x_n) + \alpha T(x_n - r_n A x_n) \\ &= [(1 - \alpha)I + \alpha T](x_n - r_n A x_n) =: S(x_n - r_n A x_n). \end{aligned}$$

Algorithm (5) potentially converges in the fixed point set faster than Algorithm (4). Here, we can see that the mapping, $S := (1 - \alpha)I + \alpha T$, satisfies the *strong*

nonexpansivity condition [11], which is a weaker condition than firm nonexpansivity. This implies that the previous algorithms in [20, 22], which can be applied to Problem (2) when T is firmly nonexpansive, cannot solve Problem (2) when T is strongly nonexpansive.

In this paper, we present an iterative algorithm for solving the variational inequality problem with a monotone, hemicontinuous operator over the fixed point set of a strongly nonexpansive mapping and show that the algorithm weakly converges to a solution to the problem under certain assumptions.

The rest of the paper is organized as follows. Section 2 covers the mathematical preliminaries. Section 3 presents the algorithm for solving the variational inequality problem for a monotone, hemicontinuous operator over the fixed point set of a strongly nonexpansive mapping, and its convergence analyses. Section 4 provides numerical comparisons of the algorithm with the previous algorithm in [20] and shows that the algorithm converges to a solution to a concrete variational inequality problem faster than the previous algorithm. Section 5 concludes the paper.

2. PRELIMINARIES

Throughout this paper, we will denote the set of all positive integers by \mathbb{N} and the set of all real numbers by \mathbb{R} . Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and its induced norm $\| \cdot \|$. We denote the strong convergence and weak convergence of $\{x_n\}$ to $x \in H$ by $x_n \rightarrow x$ and $x_n \rightharpoonup x$, respectively. It is known that H satisfies the following condition, called *Opial's condition* [33]: for any $\{x_n\} \subset H$ satisfying $x_n \rightharpoonup x_0$, $\liminf_{n \rightarrow \infty} \|x_n - x_0\| < \liminf_{n \rightarrow \infty} \|x_n - y\|$ holds for all $y \in H$ with $y \neq x_0$; see also [14, 39, 40]. To prove our main theorems, we need the following lemma, which was proven in [41]; see also [5, 39, 40].

Lemma 2.1 ([41]). *Assume that $\{s_n\}$ and $\{e_n\}$ are sequences of non-negative numbers such that $s_{n+1} \leq s_n + e_n$ for all $n \in \mathbb{N}$. If $\sum_{n=1}^{\infty} e_n < \infty$, then $\lim_{n \rightarrow \infty} s_n$ exists.*

2.1. Strong nonexpansivity and fixed point set. Let T be a mapping of H into itself. We denote the fixed point set of T by $\text{Fix}(T)$; i.e., $\text{Fix}(T) = \{z \in H : Tz = z\}$. A mapping, $T : H \rightarrow H$, is said to be *nonexpansive* if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in H$. $\text{Fix}(T)$ is closed and convex when T is nonexpansive [14, 15, 39, 40]. $T : H \rightarrow H$ is said to be *strongly nonexpansive* [11] if T is nonexpansive and if, for bounded sequences $\{x_n\}, \{y_n\} \subset H$, $\|x_n - y_n\| - \|Tx_n - Ty_n\| \rightarrow 0$ implies $\|x_n - y_n - (Tx_n - Ty_n)\| \rightarrow 0$. The following properties for strongly nonexpansive mappings were shown in [11]:

- $\text{Fix}(T)$ is closed and convex when $T : H \rightarrow H$ is strongly nonexpansive because T is also nonexpansive.
- A strongly nonexpansive mapping, $T : H \rightarrow H$, with $\text{Fix}(T) \neq \emptyset$ is *asymptotically regular* [10, 14], i.e., for each $x \in H$, $\lim_{n \rightarrow \infty} \|T^n x - T^{n+1} x\| = 0$.
- If $S, T : H \rightarrow H$ are strongly nonexpansive, then ST is also strongly nonexpansive, and $\text{Fix}(ST) = \text{Fix}(S) \cap \text{Fix}(T)$ when $\text{Fix}(S) \cap \text{Fix}(T) \neq \emptyset$.
- If $S : H \rightarrow H$ is strongly nonexpansive and if $T : H \rightarrow H$ is nonexpansive, then $\alpha S + (1 - \alpha)T$ is strongly nonexpansive for $\alpha \in (0, 1)$. If $\text{Fix}(S) \cap \text{Fix}(T) \neq \emptyset$, then $\text{Fix}(\alpha S + (1 - \alpha)T) = \text{Fix}(S) \cap \text{Fix}(T)$ [1]. In particular, since the identity mapping I is strongly nonexpansive, the mapping, $U :=$

$\alpha I + (1 - \alpha)T$, is strongly nonexpansive. Such U is said to be *averaged nonexpansive*.

Example 2.1. Let $D \subset H$ be a closed convex set, which is simple in the sense that P_D can be calculated explicitly. Furthermore, let $f: H \rightarrow \mathbb{R}$ be Fréchet differentiable and $\nabla f: H \rightarrow H$ be Lipschitz continuous; i.e., there exists $L > 0$ such that $\|\nabla f(x) - \nabla f(y)\| \leq L\|x - y\|$ for all $x, y \in H$. Then, for $r \in (0, 2/L]$, $S_r := P_D(I - r\nabla f)$ is nonexpansive [2], [19, Lemma 2.1]. Define $T: H \rightarrow H$ by

$$(6) \quad T := \alpha I + (1 - \alpha)S_r \quad (\alpha \in (0, 1)).$$

Then, T is strongly nonexpansive and $\text{Fix}(T) = \{x \in D: f(x) = \min_{y \in D} f(y)\}$.

Example 2.2. Let $D_i \subset H$ ($i = 0, 1, \dots, m$) be a closed convex set, which is simple in the sense that P_{D_i} can be calculated explicitly. Define $\Phi(x) := (1/2) \sum_{i=1}^m \omega_i d(x, D_i)$ for all $x \in H$, where $\omega_i \in (0, 1)$ with $\sum_{i=1}^m \omega_i = 1$ and $d(x, D_i) := \min\{\|x - y\|: y \in D_i\}$ ($i = 1, 2, \dots, m$). Also, we define $S: H \rightarrow H$ and $T: H \rightarrow H$ as

$$(7) \quad S := P_{D_0} \left[\sum_{i=1}^m \omega_i P_{D_i} \right], T := \alpha I + (1 - \alpha)S \quad (\alpha \in (0, 1)).$$

Then, S is nonexpansive [42, Proposition 4.2] and $\text{Fix}(S) = C_\Phi := \{x \in D_0: \Phi(x) = \min_{y \in D_0} \Phi(y)\}$. Hence, T is strongly nonexpansive and $\text{Fix}(T) = C_\Phi$. C_Φ is referred to as a *generalized convex feasible set* [12, 42] and is defined as a subset of D_0 that is closest to D_1, D_2, \dots, D_m in the mean square sense. Even if $\bigcap_{i=0}^m D_i = \emptyset$, C_Φ is well defined. $C_\Phi = \bigcap_{i=0}^m D_i$ holds when $\bigcap_{i=0}^m D_i \neq \emptyset$. Accordingly, C_Φ is a generalization of $\bigcap_{i=0}^m D_i$.

A mapping $F: H \rightarrow H$ is said to be *firmly nonexpansive* [8] if $\|Fx - Fy\|^2 \leq \langle x - y, Fx - Fy \rangle$ for all $x, y \in H$ (see also [14, 15, 35]). Every firmly nonexpansive mapping F can be expressed as $F = (1/2)(I + T)$ given some nonexpansive mapping T [14, 15, 35]. Hence, the class of averaged nonexpansive mappings includes the class of firmly nonexpansive mappings.

2.2. Variational inequality. An operator, $A: H \rightarrow H$, is said to be *monotone* if $\langle x - y, Ax - Ay \rangle \geq 0$ for all $x, y \in H$. $A: H \rightarrow H$ is said to be *hemicontinuous* [39, p. 204] if, for any $x, y \in H$, the mapping $g: [0, 1] \rightarrow H$ defined by $g(t) = A(tx + (1 - t)y)$ is continuous, where H has a weak topology. Let C be a nonempty, closed convex subset of H . The *variational inequality problem* [32, 38] for a monotone operator $A: H \rightarrow H$ is as follows (see also [30, 36, 39, 40, 43, 44]).

$$\text{find } z \in C \text{ such that } \langle y - z, Az \rangle \geq 0 \text{ for all } y \in C.$$

We denote the solution set of the variational inequality problem by $\text{VI}(C, A)$. The monotonicity and hemicontinuity of A imply that $\text{VI}(C, A) = \{z \in C: \langle y - z, Az \rangle \geq 0 \text{ for all } y \in C\}$ [39, subsection 7.1]. This means that $\text{VI}(C, A)$ is closed and convex. $\text{VI}(C, A)$ is nonempty when $A: H \rightarrow H$ is monotone and hemicontinuous, and $C \subset H$ is nonempty, compact, and convex [39, Theorem 7.1.8].

Example 2.3. Let $g: H \rightarrow \mathbb{R}$ be convex and continuously Fréchet differentiable and $A := \nabla g$. Then, A is monotone and hemicontinuous.

(i) Suppose that $f: H \rightarrow \mathbb{R}$ is as in Example 2.1 and $T: H \rightarrow H$ is defined as in (6) and set $G := \{z \in D: f(z) = \min_{w \in D} f(w)\}$. Then,

$$\text{VI}(\text{Fix}(T), A) = \left\{ x \in G: g(x) = \min_{y \in G} g(y) \right\}.$$

A solution of this problem is a minimizer of g over the set of all minimizers of f over D . Therefore, the problem has a triplex structure [19, 21, 24].

(ii) Suppose that $T: H \rightarrow H$ is defined as in (7). Then,

$$\text{VI}(\text{Fix}(T), A) = \left\{ x \in C_{\Phi}: g(x) = \min_{y \in C_{\Phi}} g(y) \right\}.$$

This problem is to find a minimizer of g over the generalized convex feasible set [22, 24, 28, 29, 42].

3. OPTIMIZATION OF VARIATIONAL INEQUALITY OVER FIXED POINT SET

In this section, we present an iterative algorithm for solving the variational inequality problem for a monotone, hemicontinuous operator over the fixed point set of a strongly nonexpansive mapping and its convergence analyses. We assume that $T: H \rightarrow H$ is a strongly nonexpansive mapping with $\text{Fix}(T) \neq \emptyset$ and that $A: H \rightarrow H$ is a monotone, hemicontinuous operator.

Algorithm 3.1.

Step 0 Choose $x_1 \in H$, $r_1 \in (0, 1)$, and $\alpha_1 \in [0, 1)$ arbitrarily, and let $n := 1$.

Step 1 Given $x_n \in H$, choose $r_n \in (0, 1)$ and $\alpha_n \in [0, 1)$ and compute $x_{n+1} \in H$ as

$$y_n := T(x_n - r_n A x_n), \quad x_{n+1} := \alpha_n x_n + (1 - \alpha_n) y_n.$$

Step 2 Update $n := n + 1$, and go to Step 1.

To prove our main theorems, we need the following lemma.

Lemma 3.1. *Suppose that $\{x_n\}$ is a sequence generated by Algorithm 3.1 and that $\{Ax_n\}$ is bounded. Moreover, assume that*

$$(A) \quad \sum_{n=1}^{\infty} r_n < \infty,$$

or

$$(B) \quad \sum_{n=1}^{\infty} r_n^2 < \infty, \quad \text{VI}(\text{Fix}(T), A) \neq \emptyset, \quad \text{and the existence of an } n_0 \in \mathbb{N} \text{ satisfying } \text{VI}(\text{Fix}(T), A) \subset \Omega := \bigcap_{n=n_0}^{\infty} \{x \in \text{Fix}(T) : \langle x_n - x, Ax_n \rangle \geq 0\}.$$

Then, $\{x_n\}$ is bounded.

Proof. Put $z_n := x_n - r_n A x_n$ for all $n \in \mathbb{N}$. We first assume that condition (A) is satisfied and choose $u \in \text{Fix}(T)$ arbitrarily. Accordingly, we have that, for any $n \in \mathbb{N}$,

$$\begin{aligned} \|x_{n+1} - u\| &= \|\alpha_n x_n + (1 - \alpha_n) y_n - u\| \\ &\leq \alpha_n \|x_n - u\| + (1 - \alpha_n) \|z_n - u\| \\ (8) \quad &= \alpha_n \|x_n - u\| + (1 - \alpha_n) \|(x_n - u) - r_n A x_n\| \\ &\leq \|x_n - u\| + r_n \|A x_n\|. \end{aligned}$$

From $\sum_{n=1}^{\infty} r_n < \infty$, the boundedness of $\{A x_n\}$, and Lemma 2.1, the limit of $\{\|x_n - u\|\}$ exists for all $u \in \text{Fix}(T)$, which implies that $\{x_n\}$ is bounded.

Next, suppose that condition (B) is satisfied, and let $u \in \text{Fix}(T)$. Then, from the monotonicity of A , we find that, for any $n \in \mathbb{N}$,

$$\begin{aligned}
\|x_{n+1} - u\|^2 &= \|\alpha_n x_n + (1 - \alpha_n)y_n - u\|^2 \\
&\leq \alpha_n \|x_n - u\|^2 + (1 - \alpha_n) \|y_n - u\|^2 \\
&\leq \alpha_n \|x_n - u\|^2 + (1 - \alpha_n) \|z_n - u\|^2 \\
&= \alpha_n \|x_n - u\|^2 + (1 - \alpha_n) \|(x_n - u) - r_n Ax_n\|^2 \\
(9) \quad &= \alpha_n \|x_n - u\|^2 + (1 - \alpha_n) (\|x_n - u\|^2 + 2r_n \langle u - x_n, Ax_n \rangle \\
&\quad + r_n^2 \|Ax_n\|^2) \\
&\leq \|x_n - u\|^2 + (1 - \alpha_n) (2r_n \langle u - x_n, Ax_n \rangle + Kr_n^2) \\
&= \|x_n - u\|^2 + (1 - \alpha_n) (2r_n \langle u - x_n, Ax_n - Au \rangle \\
&\quad + 2r_n \langle u - x_n, Au \rangle + Kr_n^2) \\
&\leq \|x_n - u\|^2 + 2r_n(1 - \alpha_n) \langle u - x_n, Au \rangle + Kr_n^2,
\end{aligned}$$

where $K := \sup\{\|Ax_n\|^2 : n \in \mathbb{N}\} < \infty$. Especially in the case of $u \in \text{VI}(\text{Fix}(T), A) \subset \Omega$, it follows from condition (B) that, for any $n \geq n_0$,

$$\begin{aligned}
\|x_{n+1} - u\|^2 &\leq \|x_n - u\|^2 + 2r_n(1 - \alpha_n) \langle u - x_n, Ax_n \rangle + Kr_n^2 \\
&\leq \|x_n - u\|^2 + Kr_n^2.
\end{aligned}$$

Hence, the condition, $\sum_{n=1}^{\infty} r_n^2 < \infty$, and Lemma 2.1 guarantee that the limit of $\{\|x_n - u\|\}$ exists for all $u \in \text{VI}(\text{Fix}(T), A)$. We thus conclude that $\{x_n\}$ is bounded. \square

Now, we are in the position to perform the convergence analysis on Algorithm 3.1 under condition (A) in Lemma 3.1.

Theorem 3.1. *Let $\{x_n\}$ be a sequence generated by Algorithm 3.1 and assume that $\{Ax_n\}$ is bounded and that the sequences $\{\alpha_n\} \subset [0, 1)$ and $\{r_n\} \subset (0, 1)$ satisfy*

$$\limsup_{n \rightarrow \infty} \alpha_n < 1, \quad \sum_{n=1}^{\infty} r_n < \infty, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\|x_n - y_n\|}{r_n} = 0.$$

Then, Algorithm 3.1 converges weakly to a point in $\text{VI}(\text{Fix}(T), A)$.

Proof. Put $z_n := x_n - r_n Ax_n$ for all $n \in \mathbb{N}$. The proof consists of the following steps:

- (a) Prove that $\{x_n\}$ and $\{z_n\}$ are bounded.
- (b) Prove that $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$ and $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ hold.
- (c) Prove that $\{x_n\}$ converges weakly to a point in $\text{VI}(\text{Fix}(T), A)$.

(a) Choose $u \in \text{Fix}(T)$ arbitrarily. From the inequality, $\|z_n - u\| = \|(x_n - r_n Ax_n) - u\| \leq \|x_n - u\| + r_n \|Ax_n\|$, and Lemma 3.1, we deduce that $\{z_n\}$ is bounded.

(b) Put $c := \lim_{n \rightarrow \infty} \|x_n - u\|$ for any $u \in \text{Fix}(T)$. Then, from $\sum_{n=1}^{\infty} r_n < \infty$, for any $\varepsilon > 0$, we can choose $m \in \mathbb{N}$ such that $\| \|x_n - u\| - c \| \leq \varepsilon$, and $r_n \leq \varepsilon$ for all $n \geq m$. Also, there exists $a > 0$ such that $\alpha_n < a < 1$ for all $n \geq m$ because of $\limsup_{n \rightarrow \infty} \alpha_n < 1$. Since $y_n = (1/(1 - \alpha_n))x_{n+1} - (\alpha_n/(1 - \alpha_n))x_n$, we have

$$\|y_n - u\| \geq \frac{1}{1 - \alpha_n} \|x_{n+1} - u\| - \frac{\alpha_n}{1 - \alpha_n} \|x_n - u\|$$

for all $n \in \mathbb{N}$. We find that, for any $n \geq m$,

$$\|y_n - u\| \geq \frac{1}{1 - \alpha_n}(c - \varepsilon) - \frac{\alpha_n}{1 - \alpha_n}(c + \varepsilon) = c - \frac{1 + \alpha_n}{1 - \alpha_n}\varepsilon \geq c - \frac{1 + a}{1 - a}\varepsilon.$$

Hence, for any $u \in \text{Fix}(T)$ and for any $n \geq m$, we have

$$\begin{aligned} 0 &\leq \|z_n - u\| - \|Tz_n - Tu\| \leq \|x_n - u\| + r_n \|Ax_n\| - \|y_n - u\| \\ &\leq c + \varepsilon + \sqrt{K}\varepsilon - \left(c - \frac{1 + a}{1 - a}\varepsilon\right) = \left(\frac{2}{1 - a} + \sqrt{K}\right)\varepsilon, \end{aligned}$$

where $K = \sup\{\|Ax_n\|^2 : n \in \mathbb{N}\} < \infty$, which implies that $\lim_{n \rightarrow \infty}(\|z_n - u\| - \|Tz_n - Tu\|) = 0$. Since T is strongly nonexpansive, we get

$$(10) \quad \lim_{n \rightarrow \infty} \|(z_n - u) - (Tz_n - Tu)\| = \lim_{n \rightarrow \infty} \|z_n - Tz_n\| = \lim_{n \rightarrow \infty} \|z_n - y_n\| = 0.$$

From (10) and $\|x_n - z_n\| = r_n \|Ax_n\| \rightarrow 0$ as $n \rightarrow \infty$, we also get

$$(11) \quad \lim_{n \rightarrow \infty} \|x_n - y_n\| = 0.$$

From $\|x_n - Tx_n\| \leq \|x_n - y_n\| + \|y_n - Tx_n\| \leq \|x_n - y_n\| + \|z_n - x_n\|$, and (11), we deduce that

$$(12) \quad \lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0.$$

(c) From the boundedness of $\{x_n\}$, there exists a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that $\{x_{n_i}\}$ converges weakly to a point $v \in H$. From the nonexpansivity of T and (12), it is guaranteed that T is demiclosed (i.e., $x_n \rightharpoonup u$ and $\|x_n - Tx_n\| \rightarrow 0$ imply $u \in \text{Fix}(T)$). Hence, we have $v \in \text{Fix}(T)$. From (9), we get, for any $u \in \text{Fix}(T)$ and for any $n \in \mathbb{N}$,

$$\begin{aligned} 0 &\leq (\|x_n - u\| + \|x_{n+1} - u\|)(\|x_n - u\| - \|x_{n+1} - u\|) \\ &\quad + 2r_n(1 - \alpha_n) \langle u - x_n, Au \rangle + Kr_n^2, \end{aligned}$$

which means

$$\begin{aligned} 0 &\leq \frac{L\|x_n - x_{n+1}\|}{r_n} + 2(1 - \alpha_n) \langle u - x_n, Au \rangle + Kr_n \\ &= \frac{L(1 - \alpha_n)\|x_n - y_n\|}{r_n} + 2(1 - \alpha_n) \langle u - x_n, Au \rangle + Kr_n \\ &\leq \frac{L\|x_n - y_n\|}{r_n} + 2(1 - \alpha_n) \langle u - x_n, Au \rangle + Kr_n, \end{aligned}$$

where $L := \sup\{\|x_n - u\| + \|x_{n+1} - u\| : n \in \mathbb{N}\} < \infty$. From $\|x_n - y_n\|/r_n \rightarrow 0$, $x_n \rightharpoonup v$, $\limsup_n \alpha_n < 1$, and $r_n \rightarrow 0$, we have

$$0 \leq \langle u - v, Au \rangle \text{ for all } u \in \text{Fix}(T).$$

The monotonicity and hemicontinuity of A imply that $v \in \text{VI}(\text{Fix}(T), A)$. Finally, we show that $\{x_n\}$ converges weakly to $v \in \text{VI}(\text{Fix}(T), A)$. Assume that another subsequence $\{x_{n_j}\}$ of $\{x_n\}$ converges weakly to w . Then, from the discussion above,

we also get $w \in \text{VI}(\text{Fix}(T), A)$. If $v \neq w$, Opial's theorem [33] guarantees that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - v\| &= \lim_{i \rightarrow \infty} \|x_{n_i} - v\| \\ &< \lim_{i \rightarrow \infty} \|x_{n_i} - w\| \\ &= \lim_{n \rightarrow \infty} \|x_n - w\| \\ &= \lim_{j \rightarrow \infty} \|x_{n_j} - w\| \\ &< \lim_{j \rightarrow \infty} \|x_{n_j} - v\| \\ &= \lim_{n \rightarrow \infty} \|x_n - v\|. \end{aligned}$$

This is a contradiction. Thus, $v = w$. This implies that every subsequence of $\{x_n\}$ converges weakly to the same point in $\text{VI}(\text{Fix}(T), A)$. Therefore, $\{x_n\}$ converges weakly to $v \in \text{VI}(\text{Fix}(T), A)$. This completes the proof. \square

Remark 3.1. The numerical examples in [20, 22, 24] show that Algorithm 3.1 satisfies $\lim_{n \rightarrow \infty} \|x_n - y_n\|/r_n = 0$ when T is firmly nonexpansive and $r_n := 1/n^\alpha$ ($1 \leq \alpha < 2$). However, when $\alpha \geq 2$, there are counterexamples that do not satisfy $\lim_{n \rightarrow \infty} \|x_n - y_n\|/r_n = 0$ [20, 22, 24].

Remark 3.2. If the sequence $\{x_n\}$ satisfies the assumptions in Theorem 3.1, we need not assume that $\text{VI}(\text{Fix}(T), A) \neq \emptyset$ or that $n_0 \in \mathbb{N}$ exists such that $\text{VI}(\text{Fix}(T), A) \subset \Omega$ in condition (B) (see also [22, Remark 7(c)]).

Remark 3.3. Let us provide the sufficient condition of the boundedness of $\{Ax_n\}$. Suppose that $\text{Fix}(T)$ is bounded and A is Lipschitz continuous. Then, we can set a bounded set V with $\text{Fix}(T) \subset V$ onto which the projection can be computed within a finite number of arithmetic operations (e.g., V is a closed ball with a large enough radius). Accordingly, we can compute

$$(13) \quad x_{n+1} := P_V(\alpha_n x_n + (1 - \alpha_n)y_n) \quad (n = 1, 2, \dots)$$

instead of x_{n+1} in Algorithm 3.1. Since $\{x_n\} \subset V$ and V is bounded, $\{x_n\}$ is bounded. The Lipschitz continuity of A means that $\|Ax_n - Ax\| \leq L\|x_n - x\|$ ($x \in H$), where $L (> 0)$ is a constant, and hence, $\{Ax_n\}$ is bounded. We can prove that Algorithm 3.1 with Equation (13) and $\{\alpha_n\}$ and $\{r_n\}$ satisfying the conditions in Theorem 3.1 (or Theorem 3.2) weakly converges to a point in $\text{VI}(\text{Fix}(T), A)$ by referring to the proof of Theorem 3.1 (or Theorem 3.2).

We prove the following theorem under condition (B) in Lemma 3.1. The essential parts of a proof are similar those of Lemma 3.1 and Theorem 3.1, so we will only give an outline of the proof below.

Theorem 3.2. *Let $\{x_n\}$ be a sequence generated by Algorithm 3.1. Assume that $\{Ax_n\}$ is bounded and that $\{\alpha_n\} \subset [0, 1)$ and $\{r_n\} \subset (0, 1)$ satisfy*

$$\limsup_{n \rightarrow \infty} \alpha_n < 1, \quad \sum_{n=1}^{\infty} r_n^2 < \infty, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\|x_n - y_n\|}{r_n} = 0.$$

If $\text{VI}(\text{Fix}(T), A) \neq \emptyset$ and if there exists $n_0 \in \mathbb{N}$ such that $\text{VI}(\text{Fix}(T), A) \subset \bigcap_{n=n_0}^{\infty} \{x \in \text{Fix}(T) : \langle x_n - x, Ax_n \rangle \geq 0\}$, then the sequence $\{x_n\}$ converges weakly to a point in $\text{VI}(\text{Fix}(T), A)$.

Proof. Put $z_n := x_n - r_n Ax_n$ for all $n \in \mathbb{N}$. As in the proof of Theorem 3.1, we proceed with the following steps:

- (a) Prove that $\{x_n\}$ and $\{z_n\}$ are bounded.
- (b) Prove that $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ holds.
- (c) Prove that $\{x_n\}$ converges weakly to a point in $\text{VI}(\text{Fix}(T), A)$.

(a) From Lemma 3.1, it follows that the limit of $\{\|x_n - u\|\}$ exists for all $u \in \text{VI}(\text{Fix}(T), A)$, and hence, $\{x_n\}$ and $\{z_n\}$ are bounded.

(b) Let $u \in \text{VI}(\text{Fix}(T), A)$ and put $c := \lim_{n \rightarrow \infty} \|x_n - u\|$. Since $\sum_{n=1}^{\infty} r_n^2 < \infty$, the condition, $r_n \rightarrow 0$, holds. As in the proof of Theorem 3.1 (b), for any $\varepsilon > 0$, there exists $m \in \mathbb{N}$ such that

$$\| \|x_n - u\| - c \| \leq \varepsilon, \text{ and } \|y_n - u\| \geq c - \frac{1+a}{1-a} \varepsilon$$

for all $n \geq m$. By $\limsup_{n \rightarrow \infty} \alpha_n < 1$, there exists $a > 0$ such that $\alpha_n < a < 1$. Since the inequality, $\|z_n - u\| = \|(x_n - r_n Ax_n) - u\| \leq \|x_n - u\| + r_n \|Ax_n\|$, holds, we have

$$\begin{aligned} 0 &\leq \|z_n - u\| - \|Tx_n - Tu\| \\ &\leq \|x_n - u\| + r_n \|Ax_n\| - \|y_n - u\| \\ &\leq c + \varepsilon + \sqrt{K} \varepsilon - \left(c - \frac{1+a}{1-a} \varepsilon \right) \\ &= \left(\frac{2}{1-a} + \sqrt{K} \right) \varepsilon, \end{aligned}$$

where $K = \sup\{\|Ax_n\|^2 : n \in \mathbb{N}\} < \infty$. This implies that $\lim_{n \rightarrow \infty} (\|z_n - u\| - \|Tx_n - Tu\|) = 0$. From the strong nonexpansivity of T , we get $\lim_{n \rightarrow \infty} \|z_n - Tx_n\| = 0$. The rest of the proof is the same as the proof of Theorem 3.1 (b). Accordingly, we obtain $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$.

(c) Following the proof of Theorem 3.1 (c), there exists a subsequence $\{x_{n_i}\} \subset \{x_n\}$ such that $\{x_{n_i}\}$ converges weakly to $v \in \text{VI}(\text{Fix}(T), A)$. Assume that another subsequence $\{x_{n_j}\}$ of $\{x_n\}$ converges weakly to w . Then, we also have that $w \in \text{VI}(\text{Fix}(T), A)$. Since the limit of $\{\|x_n - u\|\}$ exists for $u \in \text{VI}(\text{Fix}(T), A)$, Opial's theorem [33] guarantees that $v = w$. This implies that every subsequence of $\{x_n\}$ converges weakly to the same point in $\text{VI}(\text{Fix}(T), A)$, and hence, $\{x_n\}$ converges weakly to $v \in \text{VI}(\text{Fix}(T), A)$. This completes the proof. \square

As we mentioned in Section 1, to solve constrained optimization problems whose feasible set is the fixed point set of a nonexpansive mapping T , Algorithm 3.1 must converge in $\text{Fix}(T)$ early in the execution. Therefore, it would be useful to use a large parameter $\alpha \in (0, 1)$ when a strongly nonexpansive mapping is represented by $(1 - \alpha)I + \alpha T$. Theorem 3.1 has the following consequences.

Corollary 3.1. *Let $T : H \rightarrow H$ be a nonexpansive mapping with $\text{Fix}(T) \neq \emptyset$ and let $A : H \rightarrow H$ be a monotone, hemicontinuous mapping. Let $\{x_n\}$ be a sequence generated by $x_1 \in H$ and*

$$(14) \quad \begin{cases} y_n = ((1 - \alpha)I + \alpha T)(x_n - r_n Ax_n), \\ x_{n+1} = \alpha_n x_n + (1 - \alpha_n) y_n \end{cases}$$

for all $n \in \mathbb{N}$, where $\{\alpha_n\} \subset [0, 1)$, $\alpha \in (0, 1)$ and $\{r_n\} \subset (0, 1)$. Assume that $\{Ax_n\}$ is a bounded sequence and that

$$\limsup_{n \rightarrow \infty} \alpha_n < 1, \quad \sum_{n=1}^{\infty} r_n < \infty, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\|x_n - y_n\|}{r_n} = 0.$$

Then, $\{x_n\}$ converges weakly to a point in $\text{VI}(\text{Fix}(T), A)$.

Proof. Since every averaged nonexpansive mapping is strongly nonexpansive and $\text{Fix}((1 - \alpha)I + \alpha T) = \text{Fix}(T)$ for $\alpha \in (0, 1)$, Theorem 3.1 implies Corollary 3.1. \square

By following the proof of Theorem 3.2 and Corollary 3.1, we get the following.

Corollary 3.2. *Let $T : H \rightarrow H$ be a nonexpansive mapping with $\text{Fix}(T) \neq \emptyset$ and let $A : H \rightarrow H$ be a monotone, hemicontinuous mapping. Let $\{x_n\}$ be a sequence in Algorithm (14). Assume that $\{Ax_n\}$ is a bounded sequence and that*

$$\limsup_{n \rightarrow \infty} \alpha_n < 1, \quad \sum_{n=1}^{\infty} r_n^2 < \infty, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\|x_n - y_n\|}{r_n} = 0.$$

If $\text{VI}(\text{Fix}(T), A) \neq \emptyset$ and if there exists $n_0 \in \mathbb{N}$ such that $\text{VI}(\text{Fix}(T), A) \subset \bigcap_{n=n_0}^{\infty} \{x \in \text{Fix}(T) : \langle x_n - x, Ax_n \rangle \geq 0\}$, then $\{x_n\}$ converges weakly to a point in $\text{VI}(\text{Fix}(T), A)$.

4. NUMERICAL EXAMPLES

Let us apply Algorithm 3.1 and the algorithm in [20] to the following variational inequality problem.

Problem 4.1. Define $f : \mathbb{R}^{1000} \rightarrow \mathbb{R}$ and C_i ($\subset \mathbb{R}^{1000}$) ($i = 1, 2$) by

$$\begin{aligned} f(x) &:= \frac{1}{2} \langle x, Qx \rangle \quad (x \in \mathbb{R}^{1000}), \\ C_i &:= \{x \in \mathbb{R}^{1000} : \langle a_i, x \rangle \leq b_i\} \quad (i = 1, 2), \end{aligned}$$

where $Q \in \mathbb{R}^{1000 \times 1000}$ is positive semidefinite, $a_i := (a_i^{(1)}, a_i^{(2)}, \dots, a_i^{(1000)}) \in \mathbb{R}^{1000}$, and $b_i \in \mathbb{R}_+$ ($i = 1, 2$). Find $z \in \text{VI}(C_1 \cap C_2, \nabla f)$.

We set Q as a diagonal matrix with diagonal components $0, 1, \dots, 999$ and choose $a_i^{(j)} \in (0, 100)$ ($i = 1, 2, j = 1, 2, \dots, 1000$) to be Mersenne Twister pseudo-random numbers given by the `random-real` function of `srfi-27`, Gauche.¹ We also set $b_1 := 5000$ and $b_2 := 4000$. The compiler used in this experiment was `gcc`.² The double-precision floating points were used for arithmetic processing of real numbers. The language was C.

In the experiment, we used the following algorithm.

$$(15) \quad \begin{cases} y_n := ((1 - \alpha)I + \alpha P_{C_1} P_{C_2}) \left(x_n - \frac{10^{-3}}{(n+1)^{1.001}} \nabla f(x_n) \right), \\ x_{n+1} := \frac{1}{2} x_n + \frac{1}{2} y_n \quad (n \in \mathbb{N}), \end{cases}$$

¹We used the Gauche scheme shell, 0.9.3.3 [utf-8, pthreads], x86.64-apple-darwin12.4.1.

²We used gcc version 4.2.1 (Based on Apple Inc. build 5658) (LLVM build 2336.11.00).

where $\alpha \in (0, 1)$. Note that the projection P_{C_i} ($i = 1, 2$) can be computed within a finite number of arithmetic operations [3, p.406] because C_i ($i = 1, 2$) is halfspace. More precisely,

$$P_{C_i}(x) = x + \frac{\min\{0, b_i - \langle a_i, x \rangle\}}{\|a_i\|^2} a_i \quad (x \in \mathbb{R}^{1000}, i = 1, 2).$$

We can see that algorithm (15) with $\alpha := 1/2$ coincides with the previous algorithm in [20]. Hence, we compare³ algorithm (15) with $\alpha := 9/10$ with algorithm (15) with $\alpha := 1/2$ and verify that algorithm (15) with $\alpha := 9/10$ converges in $C_1 \cap C_2 = \text{Fix}(P_{C_1}P_{C_2})$ faster than algorithm (15) with $\alpha := 1/2$. We selected one hundred initial points $x = x(k) \in \mathbb{R}^{1000}$ ($k = 1, 2, \dots, 100$) as pseudo-random numbers generated by the `rand` function of the C Standard Library. We executed algorithm (15) with $\alpha := 9/10$ and algorithm (15) with $\alpha := 1/2$ for these initial points. Let $\{x_n(k)\}$ be the sequence generated by $x(k)$ and algorithm (15). Here, we define

$$D_n := \frac{1}{100} \sum_{k=1}^{100} \|x_n(k) - P_{C_1}P_{C_2}(x_n(k))\| \quad (n \in \mathbb{N}).$$

The convergence of $\{D_n\}$ to 0 implies that algorithm (15) converges to a point in $C_1 \cap C_2$.

Corollary 3.1 guarantees that algorithm (15) converges to a solution to Problem 4.1 if $\{\nabla f(x_n)\}$ is bounded and if

$$(16) \quad \lim_{n \rightarrow \infty} (n+1)^{1.001} \|x_n - y_n\| = 0.$$

To verify whether algorithm (15) satisfies condition (16), we employed

$$X_n := \frac{1}{100} \sum_{k=1}^{100} (n+1)^{1.001} \|x_n(k) - y_n(k)\| \quad (n \in \mathbb{N}),$$

where $y_n(k) := ((1-\alpha)I + \alpha P_{C_1}P_{C_2})(x_n(k) - (10^{-3}/(n+1)^{1.001})\nabla f(x_n(k)))$ ($k = 1, 2, \dots, 100, n \in \mathbb{N}$). The convergence of $\{X_n\}$ to 0 implies that algorithm (15) satisfies condition (16). We also used

$$F_n := \frac{1}{100} \sum_{k=1}^{100} f(x_n(k)) \quad (n \in \mathbb{N})$$

to check that algorithm (15) is stable.

Figure 1 indicates the behaviors of D_n for algorithm (15) with $\alpha := 9/10$ and algorithm (15) with $\alpha := 1/2$. This figure shows that $\{D_n\}$ in algorithm (15) with $\alpha := 9/10$ converges to 0 faster than $\{D_n\}$ in algorithm (15) with $\alpha := 1/2$; i.e., algorithm (15) with $\alpha := 9/10$ converges in $C_1 \cap C_2$ faster than the previous algorithm in [20].

Figure 2 compares the behaviors of X_n for algorithm (15) with $\alpha := 9/10$ and algorithm (15) with $\alpha := 1/2$ and shows that the $\{X_n\}$ generated by each algorithm converges to 0; i.e., they each satisfy (16). Therefore, from Corollary 3.1, we can conclude that they can find a solution to Problem 4.1.

We can see from Figure 3 that $\{F_n\}$ generated by the two algorithms converge to the same value. Figures 1, 2, and 3 indicate that algorithm (15) with $\alpha := 9/10$ converges to a solution to Problem 4.1 faster than the previous algorithm in [20].

³For example, we set a large parameter, i.e., much more than 1/2: $\alpha = 9/10$.

This is because algorithm (15) uses a parameter ($\alpha := 9/10$) that is larger than $1/2$ and algorithm (15) with $\alpha > 1/2$ potentially converges in the constraint set $C_1 \cap C_2$ faster than the previous algorithm in [20] with $\alpha := 1/2$.




FIGURE 1. Behavior of D_n for algorithm (15) with $\alpha := 9/10$ and algorithm (15) with $\alpha := 1/2$




FIGURE 2. Behavior of X_n for algorithm (15) with $\alpha := 9/10$ and algorithm (15) with $\alpha := 1/2$




FIGURE 3. Behavior of F_n for algorithm (15) with $\alpha := 9/10$ and algorithm (15) with $\alpha := 1/2$

5. CONCLUSION

We studied a variational inequality problem for a monotone, hemicontinuous operator over the fixed point set of a strongly nonexpansive mapping in a Hilbert space and devised an iterative algorithm for solving it. Our convergence analyses guarantee that the algorithm weakly converges to a solution under certain

assumptions. We gave numerical results to support the convergence analyses on the algorithm. The results showed that the algorithm converges to a solution to a concrete variational inequality problem faster than the previous algorithm.

ACKNOWLEDGMENTS

We are sincerely grateful to Lead Guest Editor, Qamrul Hasan Ansari, of Special Issue on Variational Analysis and Fixed Point Theory: Dedicated to Professor Wataru Takahashi on the occasion of his 70th birthday, and the two anonymous referees for helping us improve the original manuscript.

REFERENCES

- [1] K. Aoyama, Y. Kimura, W. Takahashi, and M. Toyoda, *On a strongly nonexpansive sequence in Hilbert spaces*, J. Nonlinear Convex Anal. **8** (2007), 471–489.
- [2] J.-B. Baillon and G. Haddad, *Quelques propriétés des opérateurs angle-bornés et n -cycliquement monotones*, Israel J. Math. **26** (1977), 137–150 (French, with English summary).
- [3] H. H. Bauschke and J. M. Borwein, *On projection algorithms for solving convex feasibility problems*, SIAM Rev. **38** (1996), 367–426.
- [4] H. H. Bauschke, P. L. Combettes, and S. Reich, *The asymptotic behavior of the composition of two resolvents*, Nonlinear Anal. **60** (2005), 283–301.
- [5] V. Berinde, *Iterative approximation of fixed points*, Lecture Notes in Mathematics, Springer, Berlin, 2007.
- [6] F. E. Browder, *Fixed-point theorems for noncompact mappings in Hilbert space*, Proc. Nat. Acad. Sci. U.S.A. **53** (1965), 1272–1276.
- [7] ———, *Nonexpansive nonlinear operators in a Banach space*, Proc. Nat. Acad. Sci. U.S.A. **54** (1965), 1041–1044.
- [8] ———, *Convergence theorems for sequences of nonlinear operators in Banach spaces*, Math. Z. **100** (1967), 201–225.
- [9] F. E. Browder and W. V. Petryshyn, *Construction of fixed points of nonlinear mappings in Hilbert space*, J. Math. Anal. Appl. **20** (1967), 197–228.
- [10] ———, *The solution by iteration of linear functional equations in Banach spaces*, Bull. Amer. Math. Soc. **72** (1966), 566–570.
- [11] R. E. Bruck and S. Reich, *Nonexpansive projections and resolvents of accretive operators in Banach spaces*, Houston J. Math. **3** (1977), 459–470.
- [12] P. L. Combettes and P. Bondon, *Hard-constrained inconsistent signal feasibility problems*, IEEE Trans. Signal Process. **47** (1999), 2460–2468.
- [13] P. L. Combettes, *A block-iterative surrogate constraint splitting method for quadratic signal recovery*, IEEE Trans. Signal Process. **51** (2003), 1771–1782.
- [14] K. Goebel and W. A. Kirk, *Topics in metric fixed point theory*, Cambridge University Press, Cambridge, 1990.
- [15] K. Goebel and S. Reich, *Uniform convexity, hyperbolic geometry, and nonexpansive mappings*, Marcel Dekker Inc., New York, 1984.
- [16] D. Göhde, *Zum Prinzip der kontraktiven Abbildung*, Math. Nachr. **30** (1965), 251–258.
- [17] A. A. Goldstein, *Convex programming in Hilbert space*, Bull. Amer. Math. Soc. **70** (1964), 709–710.
- [18] S. Iemoto and W. Takahashi, *Strong convergence theorems by a hybrid steepest descent method for countable nonexpansive mappings in Hilbert spaces*, Sci. Math. Jpn. **69** (2009), 227–240.
- [19] H. Iiduka, *Strong convergence for an iterative method for the triple-hierarchical constrained optimization problem*, Nonlinear Anal. **71** (2009), 1292–1297.
- [20] ———, *A new iterative algorithm for the variational inequality problem over the fixed point set of a firmly nonexpansive mapping*, Optimization **59** (2010), 873–885.
- [21] ———, *Iterative algorithm for solving triple-hierarchical constrained optimization problem*, J. Optim. Theory Appl. **148** (2011), 580–592.
- [22] ———, *Fixed point optimization algorithm and its application to power control in CDMA data networks*, Math. Program. **133** (2012), 227–242.

- [23] ———, *Fixed point optimization algorithm and its application to network bandwidth allocation*, J. Comput. Appl. Math. **236** (2012), 1733–1742.
- [24] ———, *Iterative algorithm for triple-hierarchical constrained nonconvex optimization problem and its application to network bandwidth allocation*, SIAM J. Optim. **22** (2012), 862–878.
- [25] ———, *Fixed point optimization algorithms for distributed optimization in networked systems*, SIAM J. Optim. **23** (2013), 1–26.
- [26] ———, *Acceleration method for convex optimization over the fixed point set of a nonexpansive mapping*, Mathematical Programming (2014), to appear.
- [27] H. Iiduka and I. Yamada, *A use of conjugate gradient direction for the convex optimization problem over the fixed point set of a nonexpansive mapping*, SIAM J. Optim. **19** (2009), 1881–1893.
- [28] ———, *An ergodic algorithm for the power-control games for CDMA data networks*, J. Math. Model. Algor. **8** (2009), 1–18.
- [29] ———, *Computational method for solving a stochastic linear-quadratic control problem given an unsolvable stochastic algebraic Riccati equation*, SIAM J. Control Optim. **50** (2012), 2173–2192.
- [30] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1980.
- [31] W. A. Kirk, *A fixed point theorem for mappings which do not increase distances*, Amer. Math. Monthly **72** (1965), 1004–1006.
- [32] J.-L. Lions and G. Stampacchia, *Variational inequalities*, Comm. Pure Appl. Math. **20** (1967), 493–519.
- [33] Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
- [34] S. Reich, *A limit theorem for projections*, Linear Multilinear Algebra **13** (1983), 281–290.
- [35] S. Reich and D. Shoikhet, *Nonlinear semigroups, fixed points, and geometry of domains in Banach spaces*, Imperial College Press, London, 2005.
- [36] R. T. Rockafellar and R. J.-B. Wets, *Variational analysis*, Springer-Verlag, Berlin, 1998.
- [37] K. Slavakis and I. Yamada, *Robust wideband beamforming by the hybrid steepest descent method*, IEEE Trans. Signal Process. **55** (2007), 4511–4522.
- [38] G. Stampacchia, *Formes bilinéaires coercitives sur les ensembles convexes*, C. R. Acad. Sci. Paris **258** (1964), 4413–4416 (French).
- [39] W. Takahashi, *Nonlinear Functional Analysis. Fixed Point Theory and its Applications*, Yokohama Publishers, Yokohama, 2000.
- [40] ———, *Introduction to Nonlinear and Convex Analysis*, Yokohama Publishers, Yokohama, 2009.
- [41] K. K. Tan and H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl. **178** (1993), 301–308.
- [42] I. Yamada, *The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings*, their applications (Haifa, 2000), Stud. Comput. Math., vol. 8, North-Holland, Amsterdam, 2001, pp. 473–504.
- [43] E. Zeidler, *Nonlinear functional analysis and its applications. II/B*, Springer-Verlag, New York, 1990.
- [44] ———, *Nonlinear functional analysis and its applications. III*, Springer-Verlag, New York, 1985.

(S. Iemoto) CHUO UNIVERSITY SUGINAMI HIGH SCHOOL, 2-7-1 IMAGAWA, SUGINAMI-KU, TOKYO, 167-0035 JAPAN

E-mail address: iemoto@tamacc.chuo-u.ac.jp

(K. Hishinuma) DEPARTMENT OF COMPUTER SCIENCE, MEIJI UNIVERSITY, 1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI-SHI, KANAGAWA, 214-8571 JAPAN

E-mail address: kaz@cs.meiji.ac.jp

(H. Iiduka) DEPARTMENT OF COMPUTER SCIENCE, MEIJI UNIVERSITY, 1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI-SHI, KANAGAWA, 214-8571 JAPAN

E-mail address: iiduka@cs.meiji.ac.jp